

**For Research Use Only.**  
**Not for use in diagnostic procedures.**



## Anti-Atg9A pAb

<b>CODE No.</b>	PD042
<b>CLONALITY</b>	Polyclonal
<b>ISOTYPE</b>	Rabbit Ig, affinity purified
<b>QUANTITY</b>	100 µL
<b>SOURCE</b>	Purified Ig from rabbit serum
<b>IMMUNOGEN</b>	Mouse Atg9A, 506 aa-839 aa (recombinant)
<b>FORMURATION</b>	PBS containing 50% Glycerol (pH 7.2). No preservative is contained.
<b>STORAGE</b>	This antibody solution is stable for one year from the date of purchase when stored at -20°C.

### APPLICATIONS-CONFIRMED

<u>Western blotting</u>	1:500 for chemiluminescence detection system
<u>Immunoprecipitation</u>	2.5 µL/300 µL of cell extract from 3 x 10 <sup>6</sup> cells
<u>Immunocytochemistry</u>	1:400

### SPECIES CROSS REACTIVITY on WB

Species	Human	Mouse	Rat	Hamster
Cell	293T	MEF	PC12	CHO
Reactivity	+	+	+	+

**Entrez Gene ID** 79065 (Human), 245860 (Mouse), 363254 (Rat)

- REFERENCES**
- 1) Itakura, E., *et al.*, *J. Cell Sci.* **125**, 1488-1499 (2012) [IC]
  - 2) Young, A. R., *et al.*, *J. Cell Sci.* **119**, 3888-3900 (2006)
  - 3) Yamada, T., *et al.*, *J. Bio. Chem.* **280**, 18283-18290 (2005)

For more information, please visit our web site <https://ruo.mbl.co.jp/>



## **RELATED PRODUCTS**

PM036	Anti-LC3 (polyclonal) [WB, IP, IC, IHC, FCM]
M152-3	Anti-LC3 (4E12) [WB, IP, IC, FCM, EM]
M186-3	Anti-LC3 (8E10) [WB]
PD014	Anti-LC3 (polyclonal) [WB]
PD015	Anti-LC3 (polyclonal) [IC]
PM046	Anti-LC3 (polyclonal) [WB, IC]
M115-3	Anti-LC3 (51-11) [WB]
PM045	Anti-p62/SQSTM1 (polyclonal)
M162-3	Anti-p62/SQSTM1 (5F2)
M162-A48	Anti-p62/SQSTM1-Alexa Fluor <sup>®</sup> 488 (5F2)
M162-A59	Anti-p62/SQSTM1-Alexa Fluor <sup>®</sup> 594 (5F2)
M162-A64	Anti-p62/SQSTM1-Alexa Fluor <sup>®</sup> 647 (5F2)
PM066	Anti-p62 C-terminal (polyclonal)
PD017	Anti-Beclin 1 (polyclonal)
PM037	Anti-GABARAP (polyclonal)
M135-3	Anti-GABARAP (1F4)
PM038	Anti-GATE-16 (polyclonal)
PD041	Anti-Atg2A (polyclonal)
PM034	Anti-Atg3 (polyclonal)
M133-3	Anti-Atg3 (3E8)
M134-3	Anti-Atg4B (9H5)
PM050	Anti-Atg5 (polyclonal)
M153-3	Anti-Atg5 (4D3)
PM039	Anti-Atg7 (polyclonal)
PD042	Anti-Atg9A (polyclonal)
M151-3	Anti-Atg10 (5A7)
M154-3	Anti-Atg12 (6E5)
PD036	Anti-Atg13 (polyclonal)
M183-3	Anti-Atg13 (5G4)
PD026	Anti-Atg14 (polyclonal)
M184-3	Anti-Atg14 (4H8)
PM040	Anti-Atg16L (polyclonal)
M150-3	Anti-Atg16L (1F12)
M160-3	Anti-UVRAG (1H4)
PD027	Anti-Rubicon (polyclonal)
M170-3	Anti-Rubicon (1H6)
PM069	Anti-NRF2 (polyclonal)
PD037	Anti-Tel2 (polyclonal)


8485 Autophagy Ab Sampler Set  
PM036-PN Positive control for anti-LC3 antibody

WB: Western blotting  
IP: Immunoprecipitation  
IC: Immunocytochemistry  
IHC: Immunohistochemistry  
FCM: Flow cytometry  
EM: Immuno-electron microscopy

### **SDS-PAGE & Western blotting**

- 1) Wash  $1 \times 10^7$  cells 3 times with PBS and suspend them in 1 mL of Laemmli's sample buffer, then sonicate briefly (up to 20 sec.).
- 2) Centrifuge the tube at  $12,000 \times g$  for 5 min. at  $4^\circ\text{C}$  and transfer the supernatant to another tube.
- 3) Heating the samples at  $55^\circ\text{C}$  for 5 min. and centrifuge. Load 10  $\mu\text{L}$  of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (7.5% acrylamide) for electrophoresis.
- 4) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at  $1 \text{ mA}/\text{cm}^2$  for 1 hr. in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 5) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for overnight at  $4^\circ\text{C}$ .
- 6) Wash the membrane with PBS-T (0.05% Tween-20 in PBS) (5 min. x 3 times).
- 7) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATIONS** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 8) Wash the membrane with PBS-T (5 min. x 3 times).
- 9) Incubate the membrane with the 1:10,000 anti-IgG (Rabbit)-HRP (MBL; code no. 458) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 10) Wash the membrane with PBS-T (5 min. x 3 times).
- 11) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 12) Expose to an X-ray film in a dark room for 1 min. Develop the film as usual. The condition for exposure and development may vary.

(Positive controls for Western blotting; 293T, MEF, PC12, and CHO)




#### **Western blot analysis of Atg9A**

Lane 1: 293T  
Lane 2: MEF  
Lane 3: PC12  
Lane 4: CHO  
Immunoblotted with PD042

### **Immunoprecipitation**

- 1) Wash  $1 \times 10^7$  cells 2 times with PBS and resuspend them with 1 mL of ice-cold Lysis buffer (50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.05% NP-40) containing appropriate protease inhibitors, then sonicate briefly (up to 20 sec.).
- 2) Centrifuge the tube at  $12,000 \times g$  for 5 min. at  $4^\circ\text{C}$  and transfer the supernatant to another tube.
- 3) Mix 20  $\mu\text{L}$  of 50% protein A agarose beads slurry resuspended in 300  $\mu\text{L}$  of IP buffer (10 mM Tris-HCl (pH 8.0), 500 mM NaCl, 0.1% NP-40) with primary antibody as suggested in the **APPLICATIONS**. Incubate with gently agitation for 1 hr. at room temperature. (The amount of antibody will depend on the conditions.)
- 4) Wash the beads 3 times with 1 mL of IP buffer.
- 5) Add 300  $\mu\text{L}$  of cell lysate (prepared sample from step 2), then incubate with gentle agitation for 1 hr. at room temperature.
- 6) Wash the beads 5 times with 1 mL of Lysis buffer.
- 7) Resuspend the beads in 20  $\mu\text{L}$  of Laemmli's sample buffer, heat it at  $55^\circ\text{C}$  for 5 min. and centrifuge.
- 8) Load 10  $\mu\text{L}$  of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (7.5% acrylamide) for electrophoresis.
- 9) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at  $1 \text{ mA}/\text{cm}^2$  for 1 hr. in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 10) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for overnight at  $4^\circ\text{C}$ .
- 11) Wash the membrane with PBS-T (0.05% Tween-20 in PBS) (5 min.  $\times$  3 times).
- 12) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATIONS** for 1 hr. at room temperature. (The concentration of antibody will depend on the conditions.)
- 13) Wash the membrane with PBS-T (5 min.  $\times$  3 times).
- 14) Incubate the membrane with the 1:1,000 Rabbit TrueBlot<sup>®</sup> anti-Rabbit IgG-HRP (eBioscience; code no. 18-8816-33) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hr. at room temperature.
- 15) Wash the membrane with PBS-T (5 min.  $\times$  3 times).
- 16) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 min.
- 17) Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 18) Expose to an X-ray film in a dark room for 1 min. Develop the film as usual. The condition for exposure and development may vary.

(Positive control for Immunoprecipitation; 293T)




### ***Immunoprecipitation of Atg9A from 293T***

Lane 1: IP with normal rabbit IgG (PM035)  
Lane 2: IP with PD042  
Immunoblotted with PD042

### **Immunocytochemistry**

- 1) Spread the cells on a glass slide, then incubate in a CO<sub>2</sub> incubator for one night.
- 2) Remove the culture supernatant by careful aspiration.
- 3) Fix the cells by immersing the slide in 4% paraformaldehyde (PFA)/PBS for 10 min. at room temperature (20~25°C).
- 4) Wash the slide 2 times with PBS.
- 5) Immerse the slide in 100 µg/mL of Digitonin in PBS for 10 min. at room temperature.
- 6) Wash the slide 2 times with PBS.
- 7) Add 200 µL of the primary antibody diluted with PBS as suggested in the **APPLICATIONS** onto the cells and incubate for 1 hr. at room temperature. (Optimization of antibody concentration or incubation condition is recommended if necessary.)
- 8) Wash the slide 2 times with PBS.
- 9) Add 200 µL of 1:500 anti-IgG (Rabbit)-Alexa Fluor<sup>®</sup>488 (Invitrogen; code no. A11008) diluted with PBS onto the cells. Incubate for 30 min. at room temperature. Keep out light by aluminum foil.
- 10) Wash the slide 2 times with PBS.
- 11) Promptly add mounting medium onto the slide, then put a cover slip on it.

(Positive control for Immunocytochemistry; A549)



***Immunocytochemical detection of Atg9A in A549***